

Reported Communicable Disease Cases, NC, January-December 2003 (by date of report)*

| Disease | Year-to-Date (Fourth Quarter) | | | 4 th Quarter 2003 | Comments / Note |
|--------------------------------|-------------------------------|-------|----------------|---------------------------------|-----------------|
| | 2003 | 2002 | Mean (98-2002) | | |
| Brucellosis | 1 | 2 | 2 | 1 | |
| Campylobacter | 825 | 683 | 546 | 225 | |
| Chlamydia, laboratory reports | 26066 | 24738 | 22621 | 6819 | |
| Creutzfeldt-Jakob disease | 3 | 1 | - | 2 | Note 9 |
| Cryptosporidiosis | 56 | 40 | - | 22 | Note 1 & 2 |
| Cyclosporiasis | 2 | 0 | - | 0 | Note 1 & 2 |
| Dengue | 3 | 3 | 2 | 1 | |
| E. coli, Shiga toxin-producing | 37 | 59 | 131 | 12 | Note 3 & 9 |
| Ehrlichiosis, Granulocytic | 2 | 1 | - | 2 | Note 1 & 2 |
| Ehrlichiosis, Monocytic | 28 | 13 | - | 12 | Note 1 & 2 |
| Encephalitis, California group | 25 | 13 | - | 13 | Note 1 & 4 |
| Encephalitis, Eastern Equine | 1 | 0 | - | 0 | Note 1 & 4 |
| Encephalitis, West Nile Virus | 16 | 0 | - | 13 | Note 1 & 4 |
| Foodborne, C. perfringens | 2 | 1 | 10 | 0 | |
| Foodborne, other | 34 | 281 | 65 | 1 | |
| Foodborne, staphylococcal | 85 | 75 | 29 | 81 | |
| Gonorrhea | 15085 | 15353 | 17746 | 3779 | |
| Haemophilus influenzae | 41 | 33 | 33 | 6 | |
| Hepatitis A | 124 | 209 | 180 | 52 | |
| Hepatitis B, acute | 163 | 233 | 235 | 53 | |
| Hepatitis B, chronic | 1023 | 896 | 705 | 224 | |
| Hepatitis B, Perinatal | 4 | 1 | - | 3 | Note 10 |
| Hepatitis C, acute | 13 | 29 | 26 | 2 | Note 1 & 4 |
| Hemolytic Uremic Syndr. / TTP | 3 | 2 | 3 | 1 | |
| HIV/AIDS | 2100 | 1705 | 1553 | 443 | Note 5 |
| Legionellosis | 42 | 13 | 14 | 12 | |
| Leptospirosis | 1 | 0 | 1 | 0 | |
| Listeriosis | 18 | 8 | - | 3 | Note 8 |
| Lyme disease | 155 | 137 | 72 | 78 | |
| Malaria | 25 | 22 | 29 | 6 | |
| Measles | 1 | 0 | 0 | 0 | |
| Meningococcal disease | 37 | 35 | 49 | 7 | |
| Meningitis, pneumococcal | 25 | 38 | 47 | 3 | |
| Mumps | 2 | 2 | 7 | 0 | |
| Q Fever | 2 | 2 | 1 | 1 | |
| Rabies, animal | 759 | 702 | 568 | 159 | |
| Rocky Mountain Spotted Fever | 324 | 294 | 156 | 152 | |
| Salmonellosis | 1416 | 1655 | 1366 | 458 | |
| Shigellosis | 1058 | 1074 | 483 | 243 | |
| Strepto. A, invasive | 106 | 122 | 81 | 14 | Note 2 |
| Syphilis, total | 396 | 616 | 1086 | 77 | Note 6 |
| Tuberculosis | 374 | 434 | 453 | 143 | |
| Toxic Shock Syndrome (TSS) | 2 | 5 | 4 | 1 | |
| TSS, Streptococcal | 4 | 0 | 0 | 0 | |
| Tularemia | 1 | 1 | 2 | 0 | |
| Typhoid, Acute | 9 | 2 | 2 | 2 | |
| Vaccinia | 4 | - | - | 0 | Note 9 |
| Vanco. Resistant Enterococci | 563 | 531 | 366 | 120 | Note 2 |
| Vibrio, other | 6 | 11 | 10 | 2 | Note 2 |
| Vibrio vulnificus | 9 | 4 | 4 | 4 | |
| Whooping cough | 144 | 46 | 93 | 45 | |

* Preliminary data, as of 2/4/2004. Quarters are defined as 13 weeks periods. Only diseases with cases reported in the year 2003 are listed in the table.

Notes: 1. - =Not reportable in this entire time period; 2. Reportable since 8/1/1998; 3. E. coli O157:H7 became reportable 10/1/1994; 4. Became reportable as "Hepatitis C, acute" rather than the previous "Hepatitis, non A-non B," and as "arboviral encephalitis" (coded by type) rather than "Encephalitis," both changes 8/1/1998; 5. Earliest report with HIV infection or AIDS diagnosis; 6. Primary, secondary and early latent syphilis; 7. Reportable since 7/1/1997; 8. Reportable since 7/2001; 9. Reportable since 2/15/2003; and E. coli, Shiga toxin-producing replaces E. coli O157:H7; 10. Coded as such since 2002.