

Ehrlichiosis/Anaplasmosis

2008 Case Definition

Clinical presentation

A tick-borne illness characterized by acute onset of fever and one or more of the following symptoms or signs: headache, myalgia, malaise, anemia, leukopenia, thrombocytopenia, or elevated hepatic transaminases. Nausea, vomiting, or rash may be present in some cases. Intracytoplasmic bacterial aggregates (morulae) may be visible in the leukocytes of some patients.

Clinical evidence

Any reported fever and one or more of the following: headache, myalgia, anemia, leukopenia, thrombocytopenia, or any hepatic transaminase elevation.

Laboratory evidence

For the purposes of surveillance,

1. ***Ehrlichia chaffeensis* infection** (formerly included in the category Human Monocytic Ehrlichiosis [HME]):

Laboratory confirmed:

- Serological evidence of a fourfold change in immunoglobulin G (IgG)-specific antibody titer to *E. chaffeensis* antigen by indirect immunofluorescence assay (IFA) between paired serum samples (one taken in first week of illness and a second 2-4 weeks later), **or**
- Detection of *E. chaffeensis* DNA in a clinical specimen via amplification of a specific target by polymerase chain reaction (PCR) assay, **or**
- Demonstration of ehrlichial antigen in a biopsy or autopsy sample by immunohistochemical methods, **or**
- Isolation of *E. chaffeensis* from a clinical specimen in cell culture.

Laboratory supportive:

- Serological evidence of elevated IgG or IgM antibody reactive with *E. chaffeensis* antigen by IFA, enzyme-linked immunosorbent assay (ELISA), dot-ELISA, or assays in other formats (CDC uses an IFA IgG cutoff of >1:64 and does not use IgM test results independently as diagnostic support criteria.), **or**
- Identification of morulae in the cytoplasm of monocytes or macrophages by microscopic examination.

2. ***Ehrlichia ewingii* infection** (formerly included in the category Ehrlichiosis [unspecified, or other agent]):

Laboratory confirmed:

- Because the organism has never been cultured, antigens are not available. Thus, *Ehrlichia ewingii* infections may only be diagnosed by molecular detection methods: *E. ewingii* DNA detected in a clinical specimen via amplification of a specific target by polymerase chain reaction (PCR) assay.

3. ***Anaplasma phagocytophilum* infection** (formerly included in the category Human Granulocytic Ehrlichiosis [HGE]):

Laboratory confirmed:

- Serological evidence of a fourfold change in IgG-specific antibody titer to *A. phagocytophilum* antigen by indirect immunofluorescence assay (IFA) in paired serum samples (one taken in first week of illness and a second 2-4 weeks later), **or**
- Detection of *A. phagocytophilum* DNA in a clinical specimen via amplification of a specific target by polymerase chain reaction (PCR) assay, **or**

- Demonstration of anaplasma antigen in a biopsy/autopsy sample by immunohistochemical methods, **or**
- Isolation of *A. phagocytophilum* from a clinical specimen in cell culture.

Laboratory supportive:

- Serological evidence of elevated IgG or IgM antibody reactive with *A. phagocytophilum* antigen by IFA, enzyme-linked immunosorbent Assay (ELISA), dot-ELISA, or assays in other formats (CDC uses an IFA IgG cutoff of $\geq 1:64$ and does not use IgM test results independently as diagnostic support criteria.), **or**
- Identification of morulae in the cytoplasm of neutrophils or eosinophils by microscopic examination.

4. Human ehrlichiosis/anaplasmosis – undetermined:

- See case classification

Exposure

Exposure is defined as having been in potential tick habitats within the past 14 days before onset of symptoms. A history of a tick bite is not required.

Case Classification

Confirmed: A clinically compatible case (meets clinical evidence criteria) that is laboratory confirmed.

Probable: A clinically compatible case (meets clinical evidence criteria) that has supportive laboratory results. For ehrlichiosis/anaplasmosis – an undetermined case can only be classified as probable. This occurs when a case has compatible clinical criteria with laboratory evidence to support ehrlichia/anaplasma infection, but not with sufficient clarity to definitively place it in one of the categories previously described. This may include the identification of morulae in white cells by microscopic examination in the absence of other supportive laboratory results.

Suspect: A case with laboratory evidence of past or present infection but no clinical information available (e.g. a laboratory report).